

Fall 2019 Illinois School Immunization Requirements by Grade Level

Vaccine Requirement	Preschool	First Entry into school (Kindergarten or First Grade)	Other Grades	Minimum Intervals allowed between doses and other options for proof of immunity
Diphtheria, Tetanus, Pertussis (DTP/DTaP/ or Tdap, Td)	Three doses of DTP or Dtap by 1 year of age. One additional booster by 2 nd birthday	Four or more doses of DTP/DTaP with the last dose qualifying as a booster and received on or after the 4 th birthday	Students entering grade 6: one dose of Tdap	Minimum interval: Between series doses: 4 weeks (28 days) Between series and booster : 6 months
Polio	Two doses by 1 year of age One additional dose by 2 nd birthday	Four or more doses of the same type of Polio vaccine with the last dose qualifying as a booster and received on or after the 4 th birthday	No new requirements	Minimum interval between series doses: 4 weeks (28 days) For Grade K: 6 month interval between three dose series and booster; booster must be on or after 4 th birthday.
Measles, Mumps, Rubella (MMR)	One dose on or after the 1 st birthday	Two doses of MMR vaccine The 1 st dose on or after the 1 st birthday and the 2 nd dose no less than 4 weeks(28 days) later	No new requirements	Minimum interval: 4 weeks (28 days) <u>Lab evidence of measles,mumps,or rubella or certified physician verification of mumps disease (only) by date of illness</u>
Haemophilus influenzae type b (Hib)	Refer to ACIP Hib series schedule for children 24-59 month Children without series must have one dose after 15 months of age.	Not required after the 5 th birthday (60 months of age)	Not required after 5 th birthday (60 months of age)	ACIP Hib series schedule: https://www.cdc.gov/vaccines/vpd/hib/index.html
Hepatitis B	Three doses for all children. Third dose must have been administered on or after 6 months of age (168 days)	No requirements	Students entering 6 th grade: Three doses of hepatitis B vaccine administered at recommended intervals	Minimum interval between doses: 1&2: at least 4 weeks ((28 days) 2&3: at least 2 months (56 days) 1&3: at least 4 months (112 days) <u>Lab evidence of prior or current infection</u>
Varicella (Chicken Pox)	One dose on or after 1 st birthday	Two doses of varicella The 1 st dose must have been received on or after 1 st birthday, the 2 nd dose no less than four weeks (28 days) later		Minimum interval: 4 weeks (28 days) <u>Statement from physician or health care provider verifying disease history OR laboratory evidence of varicella immunity</u>
Pneumococcal Conjugate Vaccine (PCV 13)	Refer to ACIP PCV series schedule for children 24-59 months Children without series must have one dose after 24 months of age	Not required after the 5 th Birthday (60 months of age)	Not required after the 5 th Birthday (60 months of age)	ACIP PCV series schedule: https://www.cdc.gov/vaccines/hep/acip-recs/vacc-specific/pneumo.html
Meningococcal Conjugate Vaccine	No requirements	No requirements	Students entering 6 th grade: One dose of Meningococcal Conjugate vaccine (on or after the 11 th birthday)	Two doses of Meningococcal Conjugate vaccine at entry to 12 th grade (on or after the 16 th birthday)